

Composition K2O 25 %+ 45 SO3%

Due to its high potassium content it provides high amount of potassium which is needed for development of the fruits during and before harvest. It also increases quality of the fruits harvested and extends the shelf life. It is very important for the calibration (growth) and improvement of aroma and taste. It increases absorbability for all types of soil and enables obtaining high ec (electrical conductivity) values in desired quantities due to its high pH level and high SO3 (45%) content.



| PLACES OF USE AND DOSES | | |
|-------------------------|------------------------------------|-------------|
| Fruit trees | 200 cc into the 100 liter of water | 1,5 -2Lt/da |
| Citrus | 200 cc into the 100 liter of water | 1,5 -2Lt/da |
| Banana | 200 cc into the 100 liter of water | 1,5 -2Lt/da |
| Strawberry | 200 cc into the 100 liter of water | 1,5 -2Lt/da |
| Tomato, pepper | 200 cc into the 100 liter of water | 1,5 -2Lt/da |
| Vineyards | 200 cc into the 100 liter of water | 1,5 -2Lt/da |
| Aubergine | 200 cc into the 100 liter of water | 1,5 -2Lt/da |
| Cucumber, courgette | 200 cc into the 100 liter of water | 1,5 -2Lt/da |
| Melon, watermelon | 200 cc into the 100 liter of water | 1,5 -2Lt/da |
| Potato, onion | 200 cc into the 100 liter of water | 1,5 -2Lt/da |



Çağın Gıda Dan. Tarım Ürün. Oto. İtalat. İhracat Sanayi ve Ticaret Limited Şirketi-Şafak mah. 4282 sok. No:53/3 NTALYA Tel: 90 242 227 75 35 | Fax: + 90 242 226 27 33 | info@cagintarim.com | www.cagintarim.com