

Sytofix

High absorption and solubility

Guaranteed Content

Organic Matter	5%
Water Soluble Potassium Oxide (K ₂ O)	1%
Alginic Acids	0,10%
EC (ds/m)	0,11%
Ph	8,5 – 10,5

Sytofix ;induces cell division, development by triggering protein synthesis, and the size of the cells enlarge. It accelerates the transport of nutrients from the roots to the plant.

Sytofix ;increases the flow of flowers and fruits, helps the plant to shorten the nodes and to grow

Sytofix; Fruity growth and development, early on in the day, after harvest aroma and storage support the endurance

Other benefits include delayed overgrowth of the leaves as a result of stimulated metabolism and better plant development.

It increases the photosynthesis ability of the plant by binding more carbon dioxide. It increases the resistance of the plant against bad conditions such as adverse weather conditions.

Plants	Time of use and dosage 100 ml/100 Lt water
Fruit Tree's, Citrus	1. Application: Before Flower 2. Application: At the fruit orientation 3. Application :Two or three weeks later
Vineyard	50-75 ml/100 Lt water 1. Application: Before Flower 2. Application: At the fruit orientation 3. Application :Two or three weeks later
Olive	100 ml/100 Lt water 1. Application: Before Flower 2. Application: At the fruit orientation 3. Application :Two or three weeks later
Spinach, Tomato, Pepper, Eggplant, Lettuce, Onion, Garlic	50-75 ml/da-100 Lt water 1. Application: Before Flower 2. Application: At the fruit orientation 3. Application: 15 days later
Cucumber, Melon, Watermelon, Pumpkin, Cauliflower, Cabbage	50-75 ml/da- 100 Lt water 1. Application: Before Flower 2. Application: At the fruit orientation 3. Application :Two or three weeks later
Corn	50-75 ml/da 6-8 leaf period
Cereals	50-75 ml/da 2-5 leaf period
Potato	50 ml/da 1. Application : In tuber formation 2. Application: Tuber development is ongoing
Beans, Soybean, Peas, Chickpeas	50-75 ml/da Before Flower
Strawberry	50 ml/da-100 Lt water 1. Application: Before Flower 2. Application: Fruit discoloration during
Ornamental plants	75-100 ml/100 Lt water 1. Application: When the plants are 10-20 cm in 2. Application: After 3-4 weeks
Cotton	50 ml/da 4 applications per week from flower formation



Packing
500 cc - 1lt

